

The ultimate pre-emergent BARRICADE® for turf grass and garden beds




syngenta®

BARRICADE® Herbicide controls a wide range of pre-emergent grassy weeds and importantly breaks the germination cycle of *Poa annua*. You can trust BARRICADE® to last for up to 6 months so your application window is more flexible.

BARRICADE® binds high in the soil profile ensuring turf and ornamental roots below the barrier are not affected and your turf and ornamentals stay in prime condition.

What BARRICADE® offers you:

- Up to six months pre-emergent control of grassy weeds in turf including African Lovegrass, Summer Grass, Crab Grass, Crowsfoot Grass and Winter Grass
- Low odour and non-staining formulation
- Flexible application window
- Registered for use in garden beds for broadleaf and grassy weed control
- Easy to apply spray over the top of turf and ornamentals



BARRICADE® can be used on the following established turf species:

Warm Season Turf Grasses

Bahia Grass, Buffalo Grass, Carpet Grass, Common Couch, Hybrid Couch, Kikuyu, Queensland Blue Couch, Seashore Paspalum, Zoysia

Refer to the registered label for full details.

SpraySURE | Accuracy and Performance

To achieve great results, accurate application is important.

Target	Syngenta Nozzle	Optimal Water Rate	Optimal Pressure Range
Root and soil So	XC 08	> 500 L/ha	2 to 4 bar

BARRICADE® can deliver up to 6 months residual control. Positioning of applications can be carried out well in advance of weeds germinating and still deliver season long control.

Example: Spring applications for the control of summer grassy weeds like *Digitaria* (Summer Grass and Crab Grass) and *Eleusine* (Crowsfoot Grass) should ideally be applied around the end of August. Control will last up to the end of February. Autumn applications for the control of *Poa* (Winter Grass) should ideally be applied around the end of February. Control will last up to the end of August.

BARRICADE® as a pre-emergent herbicide works in the top soil layers. Application targeted towards delivering BARRICADE® close to the soil surface is best suited to ensure optimum results. BARRICADE® needs to be washed in within 7 days after application.

Target Weeds	Rate (Turf)	Rate (Garden Beds)	SpraySure
Barnyard Grass, Crab Grass, Crowsfoot Grass, Summer Grass, Vulpia Hair Grass	1 to 3 L/ha	4 to 8 L/ha	So
African Lovegrass	2 to 3 L/ha	4 to 8 L/ha	So
Paspalum	2 to 4 L/ha	4 to 8 L/ha	So
Winter Grass	1 to 4 L/ha	4 to 8 L/ha	So
Chickweed, Cudweed, Oxalis, Prairie Grass, Ryegrass, Sow Thistle, Willowherb	n/a	4 to 8 L/ha	So

Refer to the registered label for full details and instructions for use. **F** = Foliar applied **So** = Soil applied **Su** = Surfactant required

Active Ingredient

BARRICADE® contains 476.2 g/L proflamifen, a member of the dinitroaniline herbicides.

GROUP **K1** HERBICIDE

Formulation and Packaging

5 L pack size

A suspension concentrate (SC) with minimal odour and non-staining properties. BARRICADE® mixes completely with water and produces a homogenous easy to apply suspension.



www.syngentaturf.co.nz

 **Barricade**
Herbicide

syngenta

For more information visit www.syngentaturf.co.nz or contact your local Syngenta agent.

Syngenta Crop Protection Limited, Tower 2, Level 7, 110 Symonds St, Auckland. Phone: (09) 306 1500 Fax: (09) 306 1501, www.syngenta.co.nz. ©Registered trademark of a Syngenta Group Company. ™Trademark of a Syngenta Group Company. All products written in uppercase are registered trademarks of a Syngenta Group Company. ©Syngenta 2021. PLEASE READ LABEL COMPLETELY BEFORE USE. BARRICADE® approved pursuant to the HSNO Act 1996, Approval Code HSR101240. SB 21-287.